

None

Amanita atkinsoniana Coker



OVERALL: Rocky Gorge, 7/18/2014, J. Solem.
 TOP: Rocky Gorge, 7/18/2014, J. Solem.
 STALK, VEIL, and GILLS: Rocky Gorge, 7/18/2014, J. Solem.
 ROOTED BULB: Rocky Gorge, 7/18/2014, J. Solem.
 SPORES (8.6 – 9.8 X 6.7 – 7.5 µm): Rocky Gorge, 7/18/2014, J. Solem.

None

[*Amanita atkinsoniana*](#)
Amanitaceae

J	F	M	A	M	J	J	A	S	O	N	D
						+	-	-	-	-	

ID: Cap white w/ patches/ warts; hemispheric (flatter in age). Stalk w/ skirt-like fragile ring; large base; large volva w/ concentric rings and splits. Strong unpleasant odor.

Habitat: Solitary; microrhizal in conifer and hardwood forests.

Cap: 3 – 6" [8 – 15 cm]

White/cream/pale gray-brown w/ reddish-brown warts. Convex (flatter in age; may have hanging veil fragments). Flesh white, firm. Strong odor of old tennis shoes.

Gills: Cream w/ faint reddish stain. Free. Crowded.


Spores: Cream. Ellipsoid, smooth

Stalk: 4 – 8" x 0.4 – 0.8" [10 – 20.5 cm x 1 – 2 cm]

Colored like cap. Covered w/ scattered/abundant fibrils. Ring collapses against stalk. Tapers down; turnip-shaped bulb may split. Volva collapses onto bulb.

Frequency: Uncommon.

Locations: ROGOR.

Notes:  Mycobank 209902. Rocky Gorge specimen identified by examination of spores by R. Solem.

Section: *Lepidella*.

References: BRBD 104, 17. E&S124fn. M&M 44, Phi 25, Rog.