

Tawny Grisette

Amanita fulva Persoon sensu lato



SIDE: Wincopin Trail, 6/20/2015, R. Orr.
SIDE: Woodstock, PVSP, 7/1/2015, J. Solem.
SPORES (10.2-11.0 µm [round]): Woodstock, PVSP, 7/1/2015, R. Solem.
GILLS, STALK and VOLVA: Duvall Road, 6/12/2009, R. Solem.

Tawny Grisette

[Amanita fulva](#)
Amanitaceae

J	F	M	A	M	J	J	A	S	O	N	D
					++	+++	---	---+	+		

ID: Cap orange/brown, margin lighter and grooved. Stalk white; no ring; loose, sac-like volva.

Habitat: Solitary, clustered. On disturbed ground in forests.

Cap: 1.2 – 3.9" [3 – 10 cm]
Yellow-/orange-brown, lighter at margin. Sticky at first or when wet. Conical (becomes convex/nearly flat). Margin prominently grooved.


Gills: White. Free/slightly attached. Close. Broad.

Spores: *White*. Round, smooth.

Stalk: 2.8 – 5.9" x 0.2 – 0.8" [7 – 15 cm x 5 – 20 mm]
White or a few gray scales. Slight taper upward. No ring. White sac-like volva fitting loosely around stalk (sometimes discolored grayish/red-dish-brown). Hollow.

Frequency: Common.

Locations: DANPV, GONRA, MCHPV, MURHL, NOLAP, OTHER, RKBRP, ROBNC, ROGOR, TUFTS, WFRDP, WINTR, WSKPV.

Notes:  **Do not taste!** Mycobank 212682. *A. fulva* is a European name widely used for the northeastern North American species.

Section: *Vaginata*.

References: Bar 236. BBF 58key. BBH 49. Bni 37. K&M 93. Lin 536, 115, 674. M&M 32. Phi 20. Stu 4.