

Fat-footed Clitocybe

Ampulloclitocybe clavipes
(Persoon) Redhead, Lutzoni,
Moncalvo & Vilgalys



OVERALL: Rockburn Branch Park, 10/18/2010, *R. Orr*.
CAP, STALK and BASE: Scott's Cove, 10/6/2008, *J. Solem*.
FAIRY RING: Annapolis Rock, PRSP, 11/11/2015, *J. Solem*.

Fat-footed Clitocybe

[Ampulloclitocybe clavipes](#)
Hygrophoraceae

J	F	M	A	M	J	J	A	S	O	N	D
+						---	---	---	+++	++-	

ID: Cap gray-brown; convex (depressed in age). Base bulbous/club-shaped. Aromatic.

Habitat: Solitary, scattered, groups. On ground in coniferous and hardwood forests.

Cap: 0.8 – 3.5" [2 – 9 cm]

Gray-/olive-brown, margin paler (buff in age). Moist. Smooth/felty/irregular; may have matted fibers. Convex becoming flat w/ depressed disc (w/ or w/o umbo). Margin elevated and wavy in age. Flesh whitish/watery-gray, limp in wet weather, brittle in dry. Spicy odor.

Gills: Whitish, pale yellowish/cream in age. Decurrent. Close/subdistant, often forked/crossveined. Depth 2 mm. Secondary gills.

Spores: *White*.

Stalk: 1.4 – 2.4" x 0.1 – 0.6" [3.6 – 6 cm x 4 – 15 mm]

Whitish w/longitudinal olive-gray fibrils. Tapers up. Base bulbous/club-shaped, often w/ white mycelium.

Frequency: Occasional.

Locations: ANNRK, NOLAP, RKBPR, ROGOR.

Notes: Mycobank 374059. Annapolis Rock record identified by examination of spores by *R. Solem*.

Synonym: *Clitocybe clavipes*

References: Bar 253. BBF 83, 277. BBH 94. E&S 134. K&M 101. Lin 745,320. M&M 185. Phi 61. Roo 129. Stu 155.



SPORES (6.1-6.9 X 3.6-4.1 μm): Annapolis Rock, PRSP, 10/11/2015, *R. Solem*.