

None

Arcyria cinerea (Bullard) Persoon



OVERALL: Duvall Road, 6/12/2009, J. Solem.
CLOSEUP: Columbia, 8/8/2017, A. VanSchoor.
CLOSEUP: Murray Hill, 6/21/2016, J. Solem.

None

Arcyria cinerea
Arcyriaceae

J	F	M	A	M	J	J	A	S	O	N	D
-	-	-	-	-	+	+	+	+	-	-	-

ID: Body of pale gray (almost white) cylinders on blackish stalks; single to several on a common stalk.

Habitat: Clusters. On well-decayed logs, bark, leaf litter, dung.

Fruiting Body: 1/12 – 1/6" [2 – 4 mm]

Pale gray (almost white) to light brown, ochraceous, or (less common) yellowish-brown. Finger-like clusters (2 – 20) on a single stalk.

Stalk: Pale to black.

Spores: Pale gray/light yellow. Round, smooth, hyaline.


Frequency: Common.

Locations: BRNBR, DANPV, DFMWA, FARMS, GONRA, HIRIP, MPENA, MURHL, OTHER, SYRIV, TUFTS, WINTR.

Notes: Mycibank 174952. Murray Hill specimen identified by examination of spores by R. Solem.

References: Bar 40. Myx 87.

http://slimemold.uark.edu/fungi/images/NameImages/KIMFLEMING%5CArcyria_cinerea2KF.jpg



SPORES (6.0-6.9 µm [round]): Murray Hill, 6/21/2016, R. Solem