

None

Aureoboletus auriporus (Peck) Pouzar



OVERALL: Big Branch, 8/19/2011, *R. Solem*.
 PORES and STALK: Big Branch, 8/19/2011, *R. Solem*.
 TOP: Big Branch, 8/5/2009, *J. Solem*.
 TOP: Savage Park, 9/12/2018, *J. Solem*.
 PORES and STALK: Savage Park, 9/12/2018, *J. Solem*.

None

[*Aureoboletus auriporus*](#)
 Boletaceae

J	F	M	A	M	J	J	A	S	O	N	D
						---	++	--	---		

ID: Cap pinkish-cinnamon (paler in age); coated w/ mottled fibrils, moist and sticky; convex (flat in age); bleeds pink. Pores golden-yellow (duller in age), bruise olive/dull brick red. Stalk yellow/pinkish-brown; copious white mycelium at base.

Habitat: Scattered, groups. Under oaks.

Cap: 0.8 – 3.1" [2 – 8 cm]

Pinkish-cinnamon or pinkish/vinaceous-brown (fading in age). Coated w/ matted fibrils. Moist and sticky. Convex (broadly convex/nearly flat in age). Flesh white/pale yellow, bleeds pink; cuticle and tubes easily separated from flesh.

Pores: Brilliant golden-yellow (dull yellow in age). Slowly stain olive/dull brick-red. Circular (1 – 2/mm). Tubes (6 – 15 mm). Depressed at stalk.

Spores: *Olive-brown*. Fusiform/ellipsoid, smooth.

Stalk: 1.5 – 4.5" x 0.3 – 0.6" [4 – 11.5 cm x 6 – 17 mm]

Pale yellow at apex, streaked/flushed pale pinkish-brown downward. Flesh white. Solid. Copious white mycelium at base.


Frequency: Uncommon.

Locations: BIGBR, RKBRP.

Synonyms: *Boletus auriporus*. *Boletus viridiflavus*.

Notes: Mycobank 211067. Big Branch (8/5/2009) specimen identified by examination of spores by *R. Solem*.

References: BBF 337, 364. BRB1 95, 286. BRB2 62. BBH 228. Bin 45. Bni 356. BRBD 210, 49. Phi 256. Roo 325. W&L 339-340.



SPORES (11.7-15.4 X 5.1-5.6 µm): Big Branch, 8/5/2009, *R. Solem*.