

Conifer-cone Baeospora

Baeospora myosura
(Fries) Singer



FRUITING BODY (on pine cone): Rockburn Branch Park,
11/23/2010, *J. Solem*.

OVERALL: River Hill, 10/11/2015, *J. Solem*.

CAPS: Hammond Village, 11/13/2017, *J. Solem*.

GILLS, STALKS and MYCELIUM: Rockburn Branch Park,
11/1/2010, *J. Solem*.

Conifer-cone Baeospora

Baeospora myosura
Marasmiaceae

J	F	M	A	M	J	J	A	S	O	N	D
								---	++-	+++	

ID: Cap cinnamon; convex, flesh thin. Long white mycelium at base.

Habitat: Groups. On conifer cones (white pine, loblolly pine, spruce).

Cap: 1/4 – 3/4 " [5 – 20 mm]

Cinnamon/pale tan. Moist. Smooth. Convex. Flesh very thin.

Gills: White. Attached. Very crowded. Narrow.

Spores: *White*. Elliptic, smooth, hyaline.

Stalk: 0.4 – 2" x 1/32 – 1/16" [1 – 5 cm x 1 – 2 mm]

Whitish/brownish. Minutely hairy. Long white hairs (mycelium) at base. Hollow.

Frequency: Fairly common.

Locations: ANNRK, GUILP, HAMVL, RIVHL, RKBRP, SLMLP.

Notes: Mycobank 271938.

Synonym: *Collybia conigena*.

References: Bar 260. BBF 71, 275. BBH 120. Bni 174. BRBD 117, 21.
Lin 738, 9. M&M 136. Stu 156.



SPORES (3.7-4.5 x 1.9-2.5 µm):
Hammond Village, 11/13/2017,
R. Solem.