## Two-colored Bolete Red-and-Yellow Bolete

Boletus bicolor Raddi



## Var. bicolor

OVERALL: Savage Park, 7/19/2008, *J. Solem*. PORES: Rockburn Branch Park, 8/25/2010, *R. Orr*. OVERALL: Hammond Village, 8/23/2014, *J. Solem*. SPORES 9.5 – 10.5 X 4.2 – 4.8 μm): Hammond Village, 8/23/2014, *R. Solem*.

<u>Var. subreticulatus</u> OVERALL: Savage Park, 7/19/2008, *J. Solem*.

PORES: Rockburn Branch Park, 8/25/2010, R. Orr.

## Two-colored Bolete Red-and-Yellow Bolete

Boletus bicolor Boletaceae

J F M A M J J A S O N D +++++++

**ID**: Cap dark red/pink (yellow and finely cracked in age); convex (flat/irregular in age); flesh yellow, stains blue. Pores yellow (dingy in age), usually stain greenish-blue. Stalk club-shaped; yellow at top, remainder rosy-red. Use FeSO<sub>4</sub> and KOH.

Habitat: Solitary, groups. In hardwood (oak) or mixed forests.

## **Cap**: 2 – 6" [5 – 15.5 cm]

Dark red or other shades of red/pink (turns yellow in age). Dry. Velvety when young (finely cracked in age). Convex (flat/irregular in age). Flesh pale yellow, slowly/quickly staining blue.

**Pores**: Yellow (dingy yellow/olive in age). Stain greenish-blue (sometimes). Free to attached. Angular. (1 - 2/mm). Tubes (3 - 10 mm).

Spores: Olive/gray-brown. Oblong/ventricose, smooth.

Stalk: 2 - 4" x 0.4 - 1.1" [5 - 10 cm x 1 - 3 cm] Yellow at top, red/rosy red for remainder. Cylindrical/club-shaped. Flesh yellow, quickly stains blue (or not). Solid.

Frequency: Very common.

Locations: BIGBR, DANPV, GONRA, HAMVL, HENPV, HIRIP, MCHPV, MPENA, NOLAP, PIGTL, RKBRP, SAVPK, SLMLP, SYRIV, TRIRS, TUFTS, WSTRP, WINTR.

**Notes**: Mycobank 414477. Hammond Village record verified by examination of spores by R. Solem. Stalk on variety *subreticulatus* has reticulation; var. *bicolor* does not. Cap stains black w/ FeSO4. Flesh stains blue-gray/olive-green w/ FeSO<sub>4</sub> and pale yellow w/ KOH.

**References**: Bar 163. BBF 337, 364. BRB 97, 99; 287. Bin 47. KUM 113. Lae 190. Lin 566, 412. M&M 394. Phi 258.