
None

Boletus flammeus E.A. Dick & Snell





OVERALL: Pigtail, 8/29/2010, *J. Solem.* PORES and STALK: Pigtail, 8/29/2010, *J. Solem.*  None



**ID**: Deep red; convex; dry; stalk similar; dark red pores blue when bruised.

**Habitat**: Solitary, scattered, groups. On ground in coniferous or mixed woods.

## **Cap**: 2.4 – 3.8" [6 – 12 cm]

Dark red/deep wine red. Convex (broadly convex in age). Dry (finely appressed hairs when young). Disc cracked in age. Flesh yellow (quickly bluing).

**Pores**: Dark red (dingier in age), bluing when bruised. Circular (1 - 2/mm). Tubes (8 - 20 mm).

Spores: Olive-brown.

**Stalk**:  $2.4 - 3.1^{\circ} \times 0.4 - 1.1^{\circ} [6 - 8 \text{ cm} \times 1 - 3 \text{ cm}]$ Dark red, yellow at apex, bluing when bruised. Dark red reticulation over upper portion. Solid.

Frequency: Occasional.

Locations: BRNBR, PIGTL, SVMLT.

Notes: Mycobank 309682.

**Synonym**: *Boletus rubroflammeus* 

References: BBF 344, 366. BRB1 154, 309. BRB2 107.