

None

*Boletus inedulius* (Murrill) Murrill



TOP: Wincopin Trail 7/7/2013, J. Solem.

PORES and STALK: Wincopin Trail 7/7/2013, J. Solem.

SIDE: Wincopin Trail 7/7/2013, J. Solem.

None

*Boletus inedulius*  
Boletaceae

J	F	M	A	M	J	J	A	S	O	N	D
						+	-	-	-	-	

**ID:** Grayish-white tomentose cap, yellowish to pinkish reticulate stalk; pores/flesh stain blue. Use KOH.

**Habitat:** Scattered, groups. On ground in woods (oak or hemlock).

**Cap:** 1.5 – 4.8" [4 – 12 cm]

Whitish, whitish-/pinkish-gray (olivaceous-/pale grayish-brown in age). Convex (broadly convex/nearly plane in age). Margin remains inrolled/incurved w/ sterile band. Dry, tomentose (cracked in age w/ whiteish flesh showong). Flesh whitish/yellowish, stains blue. Bitter taste.

**Pores:** Pale yellow (olive-yellow in age), stain blue, depressed at stalk in age. Pores circular (1 – 3/mm). Tubes 8 – 16 mm.

**Spores:** Olive-brown

**Stalk:** 2 – 4.5" x 3/8 – 3/4" [5 – 11.5 cm x 1 – 2 cm]

Yellow/pale yellow at top, reddish-pinkish below. Stains blackish-brown near base when handled. Reticulate (at least upper part).

**Frequency:** Uncommon.

**Locations:** WINTR.

**Notes:** KOH stains cap pale orange, flesh orange. One of the two fruiting bodies was only slightly bitter. Mycobank 267858.

**References:** Bin 61. BBF 322key. BRB 122, 297. BRBD 230fn. KUM 117. Phi 264.