

None


Buglossoporus quercinus
(Schrader) Kotlába & Pouzar



TOP: Glenelg HS, 7/12/2020, E. Concari.
PORES: Glenelg HS, 7/12/2020, E. Concari.
TOP: Glenelg HS, 7/12/2020, S. Muller.
PORES: Glenelg HS, 7/12/2020, S. Muller.
OVERALL: Glenelg HS, 7/12/2020, S. Muller.

None

Buglossoporus quercinus
Meruliaceae



ID: Whitish cap w/ dense fuzz, fan-/tongue-shaped to circular; color different in age.

Habitat: Old oak (*Quercus*) trees only; mostly in wounds where heartwood is exposed.

Fruiting Body: 4.8 – 8" x 2" [12 – 20 cm x 5 cm]
Whitish, then yellow (reddish-brown in age). Dense fuzz (brown in age). Fan-/tongue shaped to circular. Bruises yellow, then grey-pink, then reddish, finally red-brown. Smell and taste acidic.

Pores: White. Circular/slightly angular.

Stalk: Absent/very short. Single point of attachment.

Spores: White.

Frequency: Uncommon.

Locations: OTHER

Notes: Mycobank 327217

Synonym: *B. pulvinus*, *Antrodia quercina*.

References: <https://www.fungusfactfriday.com/229-buglossoporus-quercinus/>, <http://fungi.myspecies.info/all-fungi/buglossoporus-quercinus>.



SPORES (6.0-7.0 x 2.4-3.2 µm):
Glenelg HS, 7/12/2020, R Solem.