

None

Callistosporium purpureomarginatum
Fatto & A. E. Bessette



SIDE: Henryton, PVSP, 9/4/2009, *J. Solem*.
GILLS and STALK: Henryton, PVSP, 9/4/2009, *J. Solem*.
CAP: Henryton, PVSP, 9/4/2009, *R. Solem*.
MATURE GILLS and STALK: Henryton, PVSP, 9/4/2009, *R. Solem*.
SIDE: Sykesville/River Road, 6/19/2011, *J. Solem*.

None

Callistosporium purpureomarginatum
Tricholomataceae

J	F	M	A	M	J	J	A	S	O	N	D
					+	-	-	-	+	+	-

ID: Cap purplish-red. Gills pinkish-/purplish-red. Metallic aftertaste.

Habitat: Solitary, scattered, groups. Well-decayed oak logs.

Cap: 0.6 – 1.5" [1.5 – 4 cm]

Purplish-red/brownish-violet when moist, dries to pinkish-tan. Bluish/brownish-purple band (1/8", 2-3 mm) at margin. Smooth. Convex (nearly flat in age). Flesh yellowish, thin. Metallic aftertaste.

Gills: Pinkish-yellow w/ purple overtones, edges reddish-purple. Attached. Close. Secondary gill, hyaline.

Spores: *White*. Elliptic, smooth, hyaline.

Stalk: 0.8 – 1.5" x 1/16 – 1/8" [2 – 4 cm x 1.5 – 4 mm]

Reddish-purple w/ yellow toward base and apex (brownish in age). Equal. Flesh yellow. Hollow. White broad mycellium.

Frequency: Uncommon.

Locations: HENPV, SYRIV.

Notes: Mycobank 436715. Sykesville/River Road specimen identified by examination of spores by R. Solem.

References: BBF 73, 275. M&M 159fn. Roo 154.



SPORES: (5.0 – 5.8 X 3.6 – 4.1 µm): Bollman Place, 9/7/2014, *R. Solem*