

None

Caloboletus inedulius (Murrill) Vizzini



TOP: Wincopin Trail 7/7/2013, J. Solem.
PORES and STALK: Wincopin Trail 7/7/2013, J. Solem.
SIDE: Wincopin Trail 7/7/2013, J. Solem.

None

Caloboletus inedulius
Boletaceae

J	F	M	A	M	J	J	A	S	O	N	D
						+	-	+	-	-	-

ID: Grayish-white tomentose cap, yellowish to pinkish reticulate stalk; pores/flesh stain blue. Use KOH.

Habitat: Scattered, groups. On ground in woods (oak or hemlock).

Cap: 1.5 – 4.8" [4 – 12 cm]

Whitish, whitish-/pinkish-gray (olivaceous-/pale grayish-brown in age). Convex (broadly convex/nearly plane in age). Margin remains inrolled/ incurved w/ sterile band. Dry, tomentose (cracked in age w/ whiteish flesh showing). Flesh whitish/yellowish, stains blue. Bitter taste.

Pores: Pale yellow (olive-yellow in age), stain blue, depressed at stalk in age. Pores circular (1 – 3/mm). Tubes 8 – 16 mm.

Spores: Olive-brown

Stalk: 2 – 4.5" x 3/8 – 3/4" [5 – 11.5 cm x 1 – 2 cm]

Yellow/pale yellow at top, reddish-pinkish below. Stains blackish-brown near base when handled. Reticulate (at least upper part).

Frequency: Uncommon.

Locations: WINTR.

Notes: KOH stains cap pale orange, flesh orange. One of the two fruiting bodies was only slightly bitter. Mycobank 267858.

Synonym: *Boletus inedulius*

References: Bin 61. BBF 322key. Bni 359. BRB1 191. BRB2 191. BRBD 230fn. K&M 117. Phi 264. Stu 271.