

Golden Chanterelle

Cantharellus roseocanus
(Redhead, Norvell & Danell)
Redhead, Norvell & Moncalvo



OVERALL: Henryton, 7/128/2010, *R. Orr*.
OVERALL: MPENA, 8/19/2010, *R. Orr*.
OVERALL: MPENA, 8/19/2010, *R. Orr*.
OVERALL: Scott's Cove, 6/27/2014, *J Solem*.
SPORES (8.1 – 9.3 X 4.3 – 5.8 µm): Scott's Cove, 6/27/2014, *R Solem*.

Golden Chanterelle

Cantharellus roseocanus
Cantharellaceae

J	F	M	A	M	J	J	A	S	O	N	D
					+	+	+	+	+	+	-

ID: Cap yellow-orange; flat (somewhat depressed in age). Blunt gill-like ridges. Use FeSO₄.

Habitat: Solitary, scattered, groups. On ground in forests.

Cap: 1 – 4" [2.5 – 10 cm]

Bright, intense yellow-orange/yellow (may fade to nearly white). Dry. Nearly smooth. Convex to nearly flat, sometimes w/ depressed disc. Margin thin, inrolled when young, uplifted and wavy in age (sometimes crimped or lobed). Flesh white. Apricot odor; sometimes w/ peppery taste.

Fertile Surface: Pale yellow/yellow/orange; slowly stains brownish (or not). Decurrent w/ forked blunt gill-like ridges, crossveins.

Spores: *Pale yellow-orange*. Elliptical, smooth.

Stalk: 1 – 4" [2.5 – 10 cm]

Pale yellow/yellow-orange. Slowly stains brownish (or not). Smooth. Variable.

Frequency: Common.

Locations: BIGBR, BRNBR, FRANF, HENPV, HIRIP, LONGC, MPENA, PATRP, PIGTL, UMDCF, WFRDP, WSTRP, WINTR.

Notes: Mycobank 200345. Scott's Cove record verified by examination of spores by R. Solem. FeSO₄ on fertile surface olive/olive-gray. Until recently, misidentified as *C. cibarius*.

References:

http://mushroomobserver.org/name/show_name/31279?q=2X9lc and Kuo are the only correct ones. All of these others still use *C. cibarius*: Bar 250. BBF 32, 36. Bin 97. KUM 131. Lin 387, 308, 427. McK 81, 7. M&M 335. Pac 154. Phi 248. Rog. Roo 126. Rus 100, 49.