

Blackfoot Polypore
Elegant Polypore

Cerioporus varius (Persoon)
I.V. Zmitrovich & A.E. Kovalenko



OVERALL: Henryton, PVSP, 8/13/2011, *R. Orr*.
STALK and PORES: Sykesville, PVSP, 5/24/2009, *J. Solem*.
TOP: Sykesville, PVSP, 5/24/2009, *J. Solem*.
SIDE: Sykesville, PVSP, 5/24/2009, *J. Solem*.
PORES: Wincopin, 8/5/2014, *J. Solem*.
SPORES (7.8 – 10.7 X 2.9-3.5 µm): Wincopin, 8/5/2014, *R. Solem*.

Blackfoot Polypore
Elegant Polypore

Cerioporus varius
Polyporaceae

J	F	M	A	M	J	J	A	S	O	N	D
+	+	++	+-+	+++	+++	+++	+++	+++	+++	+++	---+

ID: Body tan/cinnamon (pale in age); kidney-shaped/circular. Pores white/gray (reddish in age), decurrent. Stalk tan above, base/lower portion black; central/lateral.

Habitat: Scattered. On dead branches and twigs of hardwoods.

Fruiting Body: 2 – 4" [5 – 10 cm]

Cinnamon-buff/tan (whitish in age). Dry, nearly smooth. Kidney-shaped, flat/concave. Flesh (1 – 5 mm) white/pale brown. Odor sweet.

Pores: White/gray/yellowish (light reddish-brown in age). Angular (4 – 5/mm). Tubes (2 mm) descend stalk.

Stalk: 0.2 – 3" x 0.1 – 0.6" [0.5 – 7.5 cm x 1.5 – 15 mm]

Tan/black on base or lower half; smooth/fine hairs (smooth in age); central/lateral. Flesh pale brown.

Spores: *White*. Oval, smooth.

Frequency: Abundant.

Locations: ANNRK, BENBR, BIGBR, BOLPL, BRNBR, CASCT, CEDLP, DANPV, FRANF, GONRA, HENPV, HIRIP, LONGC, MANWD, MCHPV, MPENA, MTPLT, MURHL, OTHER, PATBT, PATRP, PIGTL, ROBNC, RKBRP, ROGOR, SAVPK, SLMLP, SUKBR, SYRIV, WSKPV, WSTRP, WINTR.

Synonym: *Polyporus varius*. *P. elegans*.

Notes: Mycobank 812042. Pigtail specimen identified by examination of spores by *R. Solem*.

References: Bar 146. BRBD 267,63. E&S 321. K&M 324. Lin 483, 453. Phi 300. Roo 360fn. W&L 558.