

ID: Reddish-buff club w/ wrinkled (usually heavily wrinkled) head; stem blends into fertile surface. Use KOH.

Habitat: Solitary, groups. On ground in hardwoods (hickory-oak) or pine forests.

Fruiting Body: 1.2 - 3.9" high X 0.8 - 2.4" wide $[3 - 10 \times 2 - 6 \text{ cm}]$ Fruiting body vertically wrinkled/grooved; tapes up to wrinkled (sometimes heavily wrinkled) head. Reddish-buff (cinnamon-brown in age or bruised); may have violet-brown tones. Flesh white, firm/spongy; slowly turns brownish when cut.

Stalk: Not easily distinguished from rest of fruiting body. Up to 1.2" [4 cm] may be buried. White at base w/ radiating white cords.

Spores: White. Ellipsoid, smooth, hyaline.


Frequency: Uncommon.

Locations: GONRA.

None

Notes: Mycobank 294980. KOH turns top yellow. Gorman Stream Valley NRA specimen identified by examination of spores by R. Solem.

References: BBH 354fn. Kuo.



SPORES (8.6-10.2 x 4.2-5.5 µm): Gorman Stream Valley NRA, 10/29/2018 R. Solem.







OVERALL: Gorman Stream Valley NRA, 10/29/2018, J. Solem. OVERALL: Gorman Stream Valley NRA, 10/29/2018, J. Solem. TOP: Gorman Stream Valley NRA, 10/29/2018, J. Solem. TOP: Gorman Stream Valley NRA, 10/29/2018, J. Solem.