

None

Clitocybe eccentrica [or near]



SIDE: High Ridge Park, 6/23/2009, R. Solem.

TOP: High Ridge Park, 6/21/2009, J. Solem.

None

Clitocybe eccentrica [or near]
Tricholomataceae

| | | | | | | | | | | | |
|---|---|---|---|---|---|-----|---|-----|-----|-----|-----|
| J | F | M | A | M | J | J | A | S | O | N | D |
| | | | | | + | +++ | + | --- | --- | --- | --- |

ID: Caps variable w/ age (whitish/yellowish/brownish); convex (funnel-shaped in age). Stalk w/ white rhizomorphs at base. Strong odor.

Habitat: Groups, clusters. On wood (primarily hardwood) or woody debris.

Cap: 0.4 – 2.4" [1 – 6 cm]
Whitish/pale brownish/orangish-yellow; changes color markedly in age. Dry. Smooth. Convex (flat/shallowly funnel-shaped in age). Margin frequently inrolled and split in age. Flesh pale yellow. Strong odor.

Gills: White/buff. Decurrent. Fairly crowded.

Spores: White/creamy.

Stalk: 1.2 – 2" x 0.3" [3 – 5 cm x 7 mm]
Light yellow. Dry. Smooth/ finely hairy above; densely hairy at base. More or less equal. White rhizomorphs at base. Yellow flesh. Stuffed.

Frequency: Fairly common.

Locations: HIRIP, HTWMA, LONGC, MPENA, NOLAP, ROGOR, SAVPK, SYRIV.

Notes: *C. eccentrica* may not be exact identification; see Kuo for details.

References: Kuo.