Clitopilus prunulus (Scopoli) P. Kummer



Clitopiilus prunulus Entalomataceae












TOP: Big Branch, 8/19/2011, *R. Solem.*GILLS and STALK: Big Branch, 8/19/2011, *R. Solem.*OVERALL: Long Corner, PRST, 8/7/2009, *R. Solem.*SIDE: Long Corner, PRST, 8/7/2009, *R. Solem.*SPORES (9.0-10.6 x 5.5-5.9 µm): Big Branch, 8/19/2011, *R. Solem.*



ID: Cap and stalk gray-white. Margin inrolled. Smells like bread.

Habitat: Scattered, groups. On ground in open woods.

Cap: 1.2 - 4" [3 – 10 cm]

Whitishy/grayish. Convex/flat (sometimes sunken in age). Dry, felty. Margin inrolled (often wavy/ irregular in age). Flesh white, firm. Smells like fresh bread or bread dough.

Gills: White (pinkish in age). Decurrent. Close.

Spores: Salmon-pink. Ellipsoid w/ longitudinal ridges.

Stalk: 1 - 3" x 1/8 - 5/8" [2.5 - 7.5 cm x 3 - 15 mm]

Dull white/pale gray. Dry.

Frequency: Fairly common.

Locations: BIGBR, HAMVL, HENPV, LONGC, SLMLP, WSTRP, WINTR.

Notes: Mycobank 237006. The Big Branch and Long Corner specimens were identified by examination of spores by R. Solem.

References: Bar 189. BBF 88, 278. BBH 141. Bni 231. Lin 641, 242.

M&M 213. Phi 153. Roo 131. Stu 225.