Polyporus brumalis (Persoon) Fries

Winter Polypore

Polyporus brumalis
Polyporaceae









OVERALL: Wincopin Trails, 11/20/2010, *J. Solem.* PORES: Wincopin Trails, 11/20/2010, *J. Solem.* TOP: Wincopin Trails, 11/20/2010, *J. Solem.* TOP: Gorman Stream Valley NRA, 3/16/2011, *J. Solem.*

J F M A M J J A S O N D

ID: Overwinters. Scattered on dead hardwoods. Yellow-brown convex/depressed cap; short stalk (usually).

Habitat: Scattered. Dead wood of fallen hardwoods.

Fruiting Body: 0.6 - 2" [1.5 - 10 cm]

Yellow-/reddish-/blackish-brown. Convex/depressed. Hairy (nearly

smooth in age). Flesh thin (1 - 2 mm).

Pores: Whitish. Circular/angular (2 - 3/mm). Tubes (1 - 3 mm). Slightly decurrent.

Stalk: 0.8 - 2.4° x 1/16 - 1/4° [2 - 6 cm x 1 - 6 mm]

Grayish/brownish. Smooth/minutely hairy. Central or eccentric. Tapers up.

Lighter near apex.

Spores: White.

Frequency: Occasional.

Locations: GONRA, MPENA, WINTR.

Notes: Mycobank 543133.

References: Bar 146. BBF 384fn. Bin 405. Lin 479, 375. M&M 418fn.

Phi 300. Roo 361. Stu 348.