## Coprinellus disseminatus (Persoon) J.E. Lange



OVERALL: Henryton, PVSP, 8/13/2011, *R. Solem.* SIDE: Hammond Village, 9/13/2020, *J. Solem.* OVERALL: MPENA, 10/16/2013, *J. Solem.* GILLS, STALK: Hammond Village, 9/13/2020, *J. Solem.* CAP: Hammond Village, 9/13/2020, *J. Solem.* 



**ID**: Caps small, yellowish/brownish-gray; conical, pleated. Gills become black but do not deliquesce.

Habitat: Dense groups. On hardwood debris, bark, or buried wood.

**Cap**: 0.2 – 0.6" [0.5 – 1.5 cm]

Pale yellowish, brownish-gray (disc yellow-brown ). Minute shiny particles (smooth in age). Hemispheric/bell-shaped/conical, deeply pleated.

Gills: White, ashy/black in age. Attached. Nearly distant. Broad.

Spores: Blackish. Ellipsoid, smooth.

**Stalk**: 0.8 - 1.2" x 1/64 - 1/32" [2 - 3 cm x 0.5 - 1 mm] White/light tan/gray. Smooth/minutely hairy. Hollow. Fragile.

Frequency: Common.

Locations: BIGBR, CASFD, DANPV, GONRA, HAMVL, HENPV, MPENA, MURHL, OTHER, WSTRP, WINTR.

**Notes**: Mycobank 107842. Does not deliquesce. MPENA specimen identified by examination of spores by R. Solem.

Synonym: Coprinus disseminatus Pseudocoprinus disseminatus.

**References**: Bar 198. BBH 179. BRBD 124, 23. E&S 148. K&M 147. Lin 598, 4. M&M 227. Phi 234. Roo 155. W&L 276-7.



SPORES (7.2 - 8.7 X 5.0 - 5.7 µm): MPENA, 10/16/2013, R. Solem.