

None

Crepidotus crocophyllus (Berkeley) Saccardo



OVERALL: MPENA, 9/18/2009, R. Solem.
 GILLS: MPENA, 9/18/2009, R. Solem.

None

[*Crepidotus crocophyllus*](#) [or near]
Inocybaceae

J	F	M	A	M	J	J	A	S	O	N	D
		---	---	---	+++	+++	+++	+++	---	---	---

ID: Cap ground color buff/orange w/ dark brown fibrils. Gill color like cap.

Habitat: Scattered, clusters. On debarked hardwood or hardwood litter (rarely conifer) .

Cap: 0.6 – 2.4" [0.5 – 6 cm]

Variable (buff/yellow/orange ground color). Covered w/ dark brown fibrils. Dry. Shell-/fan-shaped, broadly convex. Flesh soft, thin.

Gills: Color variable (like cap), becoming brown from spores. Radiate from point of attachment. Crowded to close. Secondary gills.

Spores: Brown.

Stalk: None.

Frequency: Occasional.

Locations: DANPV, MANWD, MPENA, RKBRP, WSTRP.

Notes: Mycobank 187144. More than 100 species.

References: Bar 221. Bni 272fn. K&M 164, Kuo. M&M 285. Phi 243. Roo 18fn. Stu 210.