


None

Entoloma conicum (Saccardo) Hessler



GILLS and STALKS: Triadelphia Reservoir, 8/24/2012, *J. Solem*.

OVERALL: Franciscan Friars, 7/31/2013, *J. Solem*.

GILLS and STALK: Triadelphia Reservoir, 8/24/2012, *J. Solem*.

MYCELIUM: Franciscan Friars, 7/31/2013, *J. Solem*.

SPORES (10.5-10.8 X 6.2-7.6 µm): Franciscan Friars, 7/31/2013, *R. Solem*.

None

[Entoloma conicum](#)
Entalomataceae

J	F	M	A	M	J	J	A	S	O	N	D
						--+	+++	+++			

ID: Cap grayish-cinnam. Conic w/ tiny umbo.

Habitat: Scattered, groups. In moss/on humus in mixed forests.

Cap: 0.4 – 0.8" [1 – 2 cm]

Pale cinnamon (wet); grayish cinnamon (dry). Conic (broadly conic w/ tiny umbo in age). Shiny-silky (dry). Margin exceeds gills; striate (wet). Flesh brownish. Fleeting chlorine odor becomes farinaceous when crushed.

Gills: Bright flesh color (mature). Almost free. Close. Narrow.

Spores: *Salmon-pink*. Angular, 5-sided.

Stalk: 1.5 – 2.2" x 1/16" [4 – 6 cm x 1 – 2 mm]
Brown. Straight. Hollow. White mycelium.

Frequency: Fairly common.

Locations: FRANF, GONRA, PIGTL, RKBPR, ROBNC, TRIRS, UMDCF.

Notes: Mycobank 330456. Triadelphia Reservoir record identified by examination of spores by *R. Solem*.

Synonyms: *Nolanea conica*.

References: BBF 214, 303. Bni 241. Stu 238.