

Rosy Polypore

*Fomitopsis cajanderi*  
(P. Karsten) B.K. Cui, M.L. Han & Y.C. Dai



UPPER SURFACE: Savage Park, 10/27/2008, J. Solem.

OVERALL: Savage Park, 10/27/2008, J. Solem.

PORES and UPPER SURFACES: Savage Park, 10/27/2008, J. Solem.

Rosy Polypore

*Fomitopsis cajanderi*  
Fomitopsidaceae

J	F	M	A	M	J	J	A	S	O	N	D
---	---	---	---	++	++	+	+	+	+	+	---

**ID:** Body reddish-/grayish-brown; fan-shaped/irregular, often descends substrate; bumpy w/ matted hairs, margin wavy. Pores rosy-pink/pinkish-white. Pleasant odor. Use KOH.

**Habitat:** Solitary, scattered, overlapping clusters. On decaying conifers (rarely hardwoods).

**Fruiting Body:** 1 – 5.5" [2.5 – 14 cm]

Pinkish-brown/reddish-brown/grayish-brown. Corky/brittle when dry, covered w/ tiny matted hairs, smooth/roughened (but not cracked), faintly zoned. Fan-shaped/semicircular/irregular, convex/nearly flat. Margin sharp, often wavy, whitish when young. Flesh rosy-pink, stains pinkish-brown. Pleasant odor.

**Pores:** White/pinkish-yellow. Stain reddish-brown in age. Circular w/ distinct cylinders (1 – 3/mm). Tubes (1 mm). Surface often irregular and descending substrate.

**Spores:** *Whitish*.

**Frequency:** Fairly common.

**Locations:** GONRA, HENPV, RKBPR, ROBNC, SAVPK, SLMLP, TRIRS.

**Notes:** Mycobank 297484. KOH turns cap and flesh black.

**Synonym:** *Rhodofomes cajanderi* .

**References:** Bar 142. BBF 389, 396. BBH 282. Bni 447. Lin 458, 524. Phi 309. Roo 373fn. W&L 534.