

Red-belted Polypore

Fomitopsis mounceae
J.E. Haight & Nakasone



OVERALL: Rockburn Branch Park, 8/1/2010, R. Orr.
PORES: Rockburn Branch Park, 8/1/2010, R. Solem.
PORES: Rockburn Branch Park, 8/1/2010, J. Solem.
OVERALL: Rockburn Branch Park, 8/1/2010, J. Solem.

Red-belted Polypore

Fomitopsis mounceae
Fomitopsidaceae

J	F	M	A	M	J	J	A	S	O	N	D
---	---	+	---	---	+	+	+	---	---	+	+

ID: Top various colors (greenish/brownish/blackish but often reddish near margin).

Habitat: Solitary, groups. On living conifers and hardwoods.

Fruiting Body: 2.8 – 15" [7 – 38 cm]

Variouly colored (greenish-gray/grayish-brown/blackish-brown). Shelf-/hoof-shaped. Concentrically furrowed (perennial). Hard. Finely cracked/smooth. Typically reddish near margin; margin creamy yellow, thick, rounded, covered w/ sticky reddish-brown resin. Flesh creamy-white/pale brown (<1.5" [4 cm]), corky/woody.

Pores: Creamy-white, bruises yellow. Circular. (3 – 5/mm).

Spores: Whitish.

Frequency: Occasional.

Locations: HENPV, MCHPV, RKBRP.

Notes: Mycobank 826717. Similar species, *F. ochracea*, never has a red band.

Synonym: *Fomitopsis pinicola*.

References: https://www.fs.fed.us/nrs/pubs/jrnl/2019/nrs_2019_haight-j_001.pdf These for *F. pinicola*: Bar 142. BBF 388, 11. BBH 264fn. Bri 446. K&M 178. Lin 459, 531. Phi 309. Roo 373. Stu 334.