

Lacquered Bracket

Ganoderma resinaceum Boud.



OVERALL: Elkridge, 10/5/2013, J. Wilkinson.
 OVERALL: Mission Road, 6/8/2013, N. Magnusson.
 OVERALL: Jackson Pond, 8/5/2008, J. Solem.

Lacquered Bracket

[*Ganoderma resinaceum*](#)
 Ganodermataceae

J	F	M	A	M	J	J	A	S	O	N	D
		- - -	- - -	- - -	+ - -	- - -	+ - -	- - -	+ - -	- - -	- - -

ID: Body brownish-red, sticky when young, looks lacquered when old; shelf-like/irregular, concentrically zoned, margin yellowish/white. Pores white/cream (ochre in age). Stalk short, stout.

Habitat: Solitary, groups. On decaying hardwoods (especially oak).

Fruiting Body: 2 – 11.8" x up to 1.1" thick [5 – 30 cm x 3 cm]
 Brownish-red in age, yellowish/whitish near margin. Lacquered surface (sticky when young), concentrically zoned, shallowly furrowed. Semicircular/shelf-like. Flesh whitish (soft and corky), (brown in age and fibrous-tough).

Pores: White/cream (ochre in age). Bruise brown when fresh. Circular (2 – 3/mm). Tubes (5 – 20 mm).

Stalk: 1.1 x 1.5" [3 cm x 4 cm]
 Sessile or short/rudimentary, brownish-red/mahogany/blackish-brown, shiny, lacquered surface.

Spores: Brown.

Frequency: Occasional.

Locations: JACPD, OTHER.

Notes: Mycobank 123196. *Ganoderma tsugae* grows on rotting coniferous wood. Previously misidentified as *G. lucidum*.. Also see *G. curtisii*.

References: Rog