

None

Gloeoporus dichrous (Fries) Bresàdola



TOP: Wincopin Trail, 9/30/2011, J. Solem.

PORES: Wincopin Trail, 9/30/2011, J. Solem.

SPLIT TOP (showing stretched gummy layer): MPEA, 8/29/2012, J. Solem.

None

Gloeoporus dichrous
Meruliaceae

J	F	M	A	M	J	J	A	S	O	N	D
		+	-	-	-	+	+	+	+	+	-

ID: Shells w/ banded to rubbery reddish-brown fertile surface. Pores separate from cap flesh. Strong sweet odor.

Habitat: Solitary, groups, overlapping clusters. On decaying hardwoods; occasionally also on deadwood of conifers and dead polypores.

Fruiting Body: 4 x 1.6" x 0. 2" thick [10 cm x 4 cm x 5 mm] White/cream. Felty (rough/tufted/smooth in age depending on weather during growth). Margin undulating. Forms elongated shelves. Flesh white [4 mm], cottony/loose.

Pores: Pale light reddish-brown (purplish/brownish in age). Angular/round [4 – 6/mm]. Tubes rubbery/gelatinous when fresh (resinous/horny in age); tubes separate from cap flesh. When fresh, splitting the fruiting body leaves a gummy layer that can be stretched!

Spores: White.

Frequency: Common.

Locations: CEDLP, FRANF, GONRA, MCHPV, MPENA, MURHL, PIGTL, RKBRN, SLMLP, SYRIV, TRIRS, WINTR, WSTRP.

Notes: Mycobank 228918.

Synonym: *Caloporus dichrous*. *Gelatoporia dichroa*.

References: BBF 392n. BBH 266. BRBD 257, 60. K&M 184. Kuo. 268. Lin 449, 590. Phi 318. Roo 372.