

Leaflike Oyster

*Hohenbuehelia petaloides*  
(Bullard) Schulzer



TOP: Scott's Cove, 9/17/2017, *J. Solem*.  
 TOP: Scott's Cove, 9/19/2009, *J. Solem*.  
 GILLS: Scotts Cove, 9/19/2009, *J. Solem*.  
 OVERALL: Scott's Cove, 9/19/2009, *R. Solem*.  
 OVERALL: MPENA, 6/13/2012, *J. Solem*.  
 GILLS: Scott's Cove, 9/17/2017, *J. Solem*.

Leaflike Oyster

[\*Hohenbuehelia petaloides\*](#)  
Pleurotaceae

J	F	M	A	M	J	J	A	S	O	N	D
					+	+	-	-	-	-	+

**ID:** Cap tan/pale brown; fan-/funnel-shaped/decurent. Stubby stalk.

**Habitat:** Solitary, groups, clusters. On ground, on or near woody debris/mulch.

**Cap:** 1 – 3" [2.5 – 7.5 cm]  
 Tan/pale brown/grayish-brown. Minutely hairy to smooth. Fan-shaped/somewhat funnel-shaped. Flesh white, thin, gelatinous.

**Gills:** White to yellow-brown. Sometimes decurrent to base of stalk. Crowded. Narrow; minutely fringed edges. Depth = 2X cap flesh. Secondary gills.

**Stalk:** Lateral. Whitish, hairy. Short, stubby.

**Spores:** *White*. Ellipsoid, smooth.

**Frequency:** Occasional.

**Locations:** BRNBR, MPENA, ROGOR, SAVPK.

**Notes:** Mycobank 119416. Scott's Cove (9/17/2017) specimen identified by examination of spores by R. Solem.

**References:** BBF 125, 286. BRBD 138, 27. K&M 207. Lin 760, 433. Roo 20fn. Stu 121.



SPORES (6.6-7.7 X 4.2-5.2 μm): Scott's Cove, 9/13/2017, *R. Solem*.