


Brown-toothed Crust

Hydnochaete olivacea
(Schwein.) Banker



OVERALL: Benson Branch, 3/11/2011, R. Solem.
CLOSEUP: Benson Branch, 3/11/2011, R. Solem.

Brown-toothed Crust

Hydnochaete olivacea
Hymenochaetaceae

| | | | | | | | | | | | |
|-----|-----|----|----|-----|-----|---|-----|-----|-----|-----|-----|
| J | F | M | A | M | J | J | A | S | O | N | D |
| --- | --- | ++ | ++ | --- | --- | + | --- | --- | --- | +++ | +++ |

ID: Brown-toothed in age, pores of cruste flattened teeth. Use KOH.

Habitat: Usually fused into large expanses on dead hardwood (oak) branches or limbs still on tree.

Fruiting Body: 2 – 4" x 4 – 8" [5 – 10 cm x 10 – 20 cm]
Dull yellowish-/reddish-brown (brown in age). Pores quickly become spine-like w/ flattened teeth. Margin may be brownish-yellow (becomes brown in age). Flesh (1-3 mm) leathery, brown, tough, fibrous.

Frequency: Common.

Locations: BRNBR, FRANF, GONRA, HAMVL, MANWD, MCHPV, MCLUR, MPENA, PIGTL, RKBMP, ROGOR, SLMLP, UNDCF, WSTRP, ZIRNP.

Notes: Mycobank 439932. Turns dark brown/black w/ KOH.

Synonyms: *Hydnoporia fuscescens*. . *Irpex cinnamomeus*.

References: BBF 401key. Bin 349. Lin 438, 577. Phi 324. Rog.