

Reddish-brown Crust *Hydnoporia tabacina* (Sowerby) Spirin,
Miettinen & K.H. Larsson



OVERALL (fertile surface): Patapsco Female Institute, 7/30/2010,
R. Solem.

OVERALL (side): Patapsco Female Institute, 7/30/2010, *R. Solem*.

Reddish-brown Crust

[*Hydnoporia tabacina*](#)
Hymenochaetaceae

J	F	M	A	M	J	J	A	S	O	N	D
---	---	---	---	---	+	+	---	---	---	+	---

ID: Often a large, fused sheet. Fertile surface dull brown, velvety (fresh) or cracked (old and dry). Use KOH.

Habitat: Overlapping clusters on decaying hardwoods.

Fruiting Body: 0.4 – 0.8" W x 1/4 – 3/8" H [1 – 2 cm x 6 – 10 mm]

Sterile surface: Yellowish-tan/dark-brown. Shelf-like stalkless caps often fused laterally and extending 8 – 12" [20 – 30 cm]. Densely matted/woolly (nearly smooth in age). Margin whitish. Fertile surface: Dull brown w/ many dark-brown *setae*. When dry, it is uneven, smooth/finely cracked/deeply radially cracked on flattened part

Frequency: Occasional.

Locations: PAFEI, WSTRP, WINTR.

Notes: Mycobank 830599. Flesh dull brown, tough, fibrous; turns dark brown w/ KOH.

Synonym: *Pseudochaete tabacina*. *Hymenochaete tabacina*.

Reference: BBF 441, 15. Lin 439, 578