Gymnopus foetidus (Sowerby) P.M. Kirk



Gymnopus foetidus Omphalotaceae









OVERALL: Murray Hill, 6/6/2016, *J. Solem.*TOPS: Murray Hill, 6/6/2016, *J. Solem.*TOPS: Schooley Mill Park, 8/1/2018, *J. Solem.*GILLS and STALKS: Murray Hill, 6/6/2016, *J. Solem.*GILLS and STALK: Schooley Mill Park, 8/1/2018, *J. Solem.* 

J F M A M J J A S O N D +++-+-+-+

**ID**: Thin, reddish-brown pleated cap in clusters.

Habitat: Groups. On hardwood limbs, logs, and litter.

**Cap**: 0.4 - 1.1" [1 - 3 cm]

Reddish-brown (tan in age). Dry. Thin. Convex/nearly flat w/ radial pleats and tiny navel-like center.

Gills: Yellowish (somewhat pinkish-red in age). Attached to collarium.

**Spores**: White. Ellipsoid/teardrop-shaped, smooth, hyaline.

**Stalk**: 0.6 – 1" X 1/16" [1.5 – 2.5 cm X 1 – 1.5 mm]

Brown. Velvety (white hairs). Tough.

Frequency: Uncommon.

Locations: MURHL.

**Notes**: Mycobank 550462. Murray Hill and Schooley Mill Park specimens identified by examination of spores by R. Solem. Murray Hill fruiting boties were on several year-old logs with bark that were supporting a footbridge. Schooley Mill Park fruiting bodies were attached to mixed wood mulch (not conifer needles).

**Synonyms**: Marasmius foetidus, Micromphale foetidum.

References: Bar 261. BBF 197key. Lin 778, 28. M&M 189.

 $\underline{https://it.wikipedia.org/wiki/Gymnopus\_foetidus}$ 



SPORES (5.7-7.3 x 2.7-3.3 µm): Schooley Mill Park, 8/1/2018, R. Solem.