

None

*Hygrocybe reidii* Kühner



OVERALL: Bollman Place, 8/6/2018, *J. Solem*.  
 OVERALL: Bollman Place, 8/6/2018, *J. Solem*.  
 CAP: Bollman Place, 8/6/2018, *J. Solem*.  
 GILLS and STALK: Bollman Place, 8/6/2018, *J. Solem*.

None

[\*Hygrocybe reidii\*](#)  
 Hygrophoraceae



J	F	M	A	M	J	J	A	S	O	N	D
					---	---	+	---	---		

**ID:** Scarlet/red-orange cap; decurrent yellow/pale orange gills.

**Habitat:** Groups. On ground in hardwood forests.

**Cap:** 0.4 – 2" [1 – 5 cm]

Orange/orange-red/scarlet (paler in age). Hemispheric (convex/nearly plane in age). Dry. Smooth when moist. Margin incurved; may be uplifted in age.

**Gills:** Dull yellow/pale orange/pinkish-orange. Decurrent. Distant. Broad. Waxy. Often w/ pale edges. Many short gills at edge.

**Spores:** *White*. Ellipsoid, smooth, hyaline.

**Stalk:** 0.8 – 2.8" x 1/16 – 1/3" [2 – 7 cm x 2 – 8 mm]


Pale orange. Smooth. Dry. Fragile. Equal/tapers down.

**Frequency:** Uncommon.

**Locations:** BOLPL.

**Notes:** Mycobank 315399. Bollman Place specimen identified by examination of spores by R. Solem.

**References:** BBF 131key. BRSB 49,113.



SPORES (6.9-8.2 x 3.5-4.6 μm):  
 Bollman Place, 8/6/2018,  
*R. Solem*.