

Caesar's Fiber Head

Inocybe heimii Bon [or near]



OVERALL: Big Branch, 9/19/2011, R. Solem.
GILLS and STALKS: Big Branch, 9/19/2011, R. Solem.

Caesar's Fiber Head

[*Inocybe heimii*](#) [or near]
Inocybaceae

J	F	M	A	M	J	J	A	S	O	N	D
						---	---	+++	---		

ID: Caps buff/tawny; densely fibrous. Stalk same.

Habitat: Solitary, scattered, groups. On ground in grass or in hardwoods.

Cap: 1 – 2" [2.5 – 5 cm]

Yellow-ochre/buff/tawny. Broadly conic/nearly flat. Densely woolly-fibrous (partly scaly in age). Dry. Margin inrolled at first. Flesh white.

Gills: Dull ochre-yellow w/ white fringes (rust-ochre in age). Attached. Close. Broad.

Spores: *Brown*. Ellipsoid, smooth.

Stalk: 0.6 – 1.6" x 1/16 – 1/4" [1.5 – 4 cm x 1.5 – 5 mm]

Yellow-ochre/tawny. Densely cottony-fibrous/scaly.

Frequency: Occasional.

Locations: BIGBR, HAMVL, HENPV, WINTR.

Notes: Mycobank 106922. Guilford Park specimen identified by examination of spores by R. Solem.

Synonym: *I. caesariata*

References: Lin 627, 314. Phi 209.