Inocybe lacera (Fries) P. Kummer

Inocybaceae

Inocybe lacera










TOP: Big Branch, 9/2/2014, J. Solem.

GILLS and STALK: Big Branch, 9/2/2014, J. Solem.

OVERALL: Big Branch, 9/5/2014, J. Solem. OVERALL: Big Branch, 9/5/2014, J. Solem.

SPORES (12.1-13.5 X 4.5-5.0 μm): Big Branch, 9/2/2014, R. Solem.



ID: Shaggy dull brown cap and stalk. Use KOH.

Habitat: Scattered, groups. Sandy soil under conifers/mixed woods.

Cap: 0.4 - 1.5" [1 - 4 cm]

None

Dull brown (tawny in age). Fibrillose to scaly. Margin often ragged/torn. Flesh whitish.

Gills: Whitish (grayish-brown in age). Attached/notched.

Spores: *Brown*. Cylindrical, smooth.

Stalk: 1.1 - 2.4" x 1/8 - 1/2" [3 - 6 cm x 3 - 6 mm]

Grayish-brown, fibrillose. Dry.

Frequency: Uncommon. **Locations**: BIGBR, PIGTL.

Notes: Mycobank 244397. Big Branch (9/2/2014 & 6/26/2016) specimens identified by examination of spores by R. Solem. KOH turns cap gray.

References: BBF 150, 291. BBH176. K&M 222. Kuo. Lin 630, 315.

M&M 303. Phi 211.