

Resinous Polypore

Ischnoderma resinosum
(Schrader) P. Karsten



OVERALL: Schooley Mill Park, 10/21/2009, J. Solem.
OVERALL (young): Henryton, PVSP, 10/2/2009, R. Solem.
PORES: Schooley Mill Park, 10/21/2009, J. Solem.
PORES EXUDING FLUID: Pigtail, 10/8/2010 R. Orr.
OVERALL: Schooley Mill Park, 11/3/2010, R. Orr.

Resinous Polypore

[*Ischnoderma resinosum*](#)
Fomitopsidaceae

J	F	M	A	M	J	J	A	S	O	N	D
			+	-	-	-	+	-	+	+	-

ID: Body brownish-orange or darker; semicircular, often descends substrate (young are different), concentrically furrowed, margin thick and lighter. Pores white (bruise brown and in age).

Habitat: Solitary, overlapping clusters. On decaying wood (oak).

Fruiting Body: 3 – 10" [7.5 – 25.5 cm]

Dull brownish-orange to dark brown. Surface concentrically or radially furrowed (faintly/distinctly zoned in age); fleshy, soft and velvety when young, later covered w/ thin, glossy, resinous crust. Semicircular/fan-shaped/convex. Margin white/ochre, thick, rounded, exudes drops of clear/amber fluid when young. Flesh (up to 2 cm) whitish/pale yellow; soft, becoming fibrous in age.

Pores: White (pale brown in age); slowly bruising brown. Pores angular/circular (4 – 6/mm). May extend down substrate.

Spores: Whitish.

Frequency: Very common.

Locations: BIGBR, CASCT, GWACP, HENPV, LONGC, MANWD, MCHPV, MPENA, OTHER, PIGTL, RKBRP, ROBNC, SLMLP, SYRIV, UMDCF, WSTRP, WINTR.

Notes: Mycobank 100840. Globular/irregular when very young.

References: Bar 140. BBF 391, 398. BBH 268. Bin 363. Bni 448. BRBD 261, 61. K&M 225. Lin 468, 525. Phi 306. Roo 377. W&L 542-3.