

Common Laccaria
Deceiver

Laccaria laccata
(Scopoli) Cooke



SIDE: High Ridge Park, 6/18/2009, *J. Solem*.
 TOPS: River Hill, 11/14/2008, *S. Earp*.
 TOP: David Force WMA, 5/20/2009, *J. Solem*.
 GILLS and STALKS: McClures, 10/24/2014, *J. Solem*.
 OVERALL: Bollman Place, 8/24/2014, *J. Solem*.
 SPORES (7.9-8.3 X 7.6-8.2 μ m): Bollman Place,
 8/24/2014, *R. Solem*.

Common Laccaria
Deceiver

[Laccaria laccata](#)
Hydnangiaceae

J	F	M	A	M	J	J	A	S	O	N	D
			+	-	+	+	+	+	+	+	+

ID: Cap orange/red-brown; convex (slightly depressed in age). Gills pinkish/flesh color. Stalk often twisted/flattened.

Habitat: Solitary, groups, small clusters. On ground in wide variety of habitats (mown grass, mixed woods, in moss under pines).

Cap: 0.4 – 2.2" [1 – 6 cm]
Orange/red-brown (pale tan in age), fading in sunlight. Finely fibrillose/smooth, usually breaks up early into minute, very soft upturned scales. Dry. Convex/broadly convex, slightly depressed. Margin translucent/grooved (or not) w/ partial veil remnants. Flesh color like cap. Thin (1 mm).

Gills: Pinkish/flesh color (sometimes purplish in age); slightly waxy. Attached/notched/subdecurrent/decurrent. Close to distant. Broad. Thick to thin.

Spores: *White*. Round/nearly round; spines 1-2.5 μ m.

Stalk: 0.6 – 4" x 1/8 – 3/8" [1.5 – 10 cm x 3 – 10 mm]
Orangish/red-brown (buff in age). Fibrous. Longitudinally grooved (or not); often twisted/flattened. Sometimes tapered upward or w/ slight bulb. Base whitish and hairy.

Frequency: Very common.

Locations: ANNRK, BIGBR, BRNBR, DANPV, DFWMA, GONRA, HAMVL, HIRIP, HOUCH, LONGC, MCLUR, NOLAP, OTHER, PIGTL, RIVHL, ROGOR, RKBPR, SAVPK, SLMLP, WINTR, WSTRP.

Notes: Mycobank 119173. Annapolis Rock, Bollman Place, and McClure specimens identified by examination of spores by *R. Solem*.

References: Bar 279. BBH 101. BBF 155, 292. Bni 130 BRBD 146, 30. Lin 762, 335. M&M 108. Phi 69. Roo 181fn.