Pine Chicken of the Woods

Laetiporus huroniensis Burdsall & Banik



Pine Chicken of the Woods

J F M A M J J A S O N D ++----

ID: Rosette of flattened caps w/ lemon yellow pores when fresh. Cap bright orange; velvety; margin blunt; wavy, often lobed.

Habitat: Solitary, overlapping clusters or rosettes. On conifers only.

Fruiting Body: $2 - 12^{\circ}$ x up to 0.8" [5 - 31 cm x 2 cm] Bright orange, then orange-yellow; bleaches white in age. Dry. Surface velvety/densely matted and woolly; radially wrinkled and roughened. Fan-/petal-shaped. Margin pale orange, blunt, dry, often lobed. Flesh pale vellow, fleshy/firm, moist.

Pores: Bright lemon yellow, creamy yellow in age. Angular (circular when young); 2 - 4/mm, tubes 1 - 5 mm.

Spores: White.

Frequency: Uncommon.

Locations: MPENA.

Notes: Mycobank 372855.

References: M&S 33.. Stu 309. http://botit.botany.wisc.edu/toms_fungi/jul2001.html. http://www.messiah.edu/Oakes/fungi_on_wood/poroid%20fungi/species% 20pages/Laetiporus%20huroniensis.htm. https://www.fpl.fs.fed.us/documnts/pdf2001/burds01a.pdf

TOP: MPENA, 4/26/2017, *J. Solem.* PORES: MPENA, 4/26/2017, *J. Solem.*