

None

Leccinellum albellum
(Peck) Bresinsky & Manfr. Binder



OVERALL: Long Corner, PRSP, 8/7/2009, *R. Solem*.
 PORES and STALK: Long Corner, PRSP, 8/7/2009, *J. Solem*.
 SPLIT CAP: Western Regional Park, 8/9/2009, *J. Solem*.
 PORES and STALK: Western Regional Park, 8/9/2009, *J. Solem*.
 OVERALL: Western Regional Park, 8/17/2011, *J. Solem*.
 PORES and STALKS: Western Regional Park, 8/17/2011, *J. Solem*.

None

Leccinellum albellum
Boletaceae

J	F	M	A	M	J	J	A	S	O	N	D
						-	+++	+++	+++	---	---

ID: Cap color variable (whitish/yellowish/brownish/gray); pitted and cracked in age; hemispheric. Pores white (brownish in age). Stalk whitish or colored like cap, rough w/ whitish scabers.

Habitat: Solitary, groups. On ground in hardwood forests (especially oak).

Cap: 1 – 2.5" [2.5 – 6.5 cm]
 Variable (whitish/yellowish/grayish-buff/pale tan/brownish-gray). Dry. Smooth to pitted; cracked in age. Hemispheric. Flesh white.

Pores: White (yellowish-gray/brownish in age). Angular (< 1/mm). Tubes (<= 10 mm).

Spores: *Brown/olive-brown*. Fusiform, smooth.

Stalk: 1.6 – 3.5" x 0.2 – 0.5" [4 – 9 cm x 5 – 12 mm]
 Whitish or colored like cap. Surface rough (whitish/brown scabers in lines in age). Cylindrical/tapers up, often twisted. Solid/stuffed.

Frequency: Common.

Locations: BIGBR, BRNBR, HIRIP, DANPV, GONRA, LONGC, MANWD, MCHPV, MPENA, PIGTL, SAVPK, TUFTS, WSTRP, WINTR, WSKPV.

Synonym: *Leccinum albellum*.

Notes: Mycobank 487801. Western Regional Park 8/17/2011 specimen identified by examination of spores by *R. Solem*.

Reference: Bin 165. BBH 205. Bni 390. BRB1 322. BRB2 238. BRBD 233, 54. E&S 259. K&M 252. Kuo. M&M 371. Phi 279. Roo 332. W&L 249.



SPORES (14.4-16.4 X 5.7-6.2 µm):
 Western Regional Park 8/17/2011,
R. Solem.