Leccinum longicurvipes (Snell & A.H. Smith) M. Kuo & B. Ortiz









OVERALL: High Ridge Park, 9/6/2009, R. Solem.

PORES and STALK: High Ridge Park, 9/9/2009, R. Solem.

SIDE: Schooley Mill Park, 8/1/2018, J. Solem.

OVERALL: Schooley Mill Park, 8/1/2018, J. Solem. None

J F M A M J J A S O N D

ID: Cap yellow-orange (olive-green in age), sticky when fresh; convex (broader in age). Pores pale yellow. Stalk whitish/ yellowish; scurfy scales reddish/brown in age. Use KOH and NH_4OH .

Habitat: Solitary, scattered, groups. On ground in oak/pine forests.

Cap: 0.6 - 2.4" [1.6 - 6 cm]

Yellow-brown/yellow-orange/orange/reddish-orange/orange-brown (often olive-green w/ orange center in age). Smooth/slightly wrinkled/pitted; sticky/slimy when fresh. Convex (broadly convex in age). Flesh (8 mm) white, becoming pale yellowish. No change when cut.

Pores: Pale-/dingy-/olive-yellow or greenish-gray. Circular/angular (2/mm). Tubes (8 – 12 mm). Pores may stain brownish.

Spores: *Brown*. Narrowly fusiform/oblong, smooth, pale brown.

Stalk: 2 - 3.5" x 0.4 - 0.8" [5 - 9 cm x 1 - 2 cm]

Whitish/yellowish or w/ reddish-pinkish-tan tones. Scurfy w/ tiny scales yellow (reddish-brown in age). Nearly equal/tapers up; often curved above base. Solid. White mycelium.


Frequency: Fairly common.

Locations: BRNBR, HENPV, HIRIP, MANWD, NOLAP, PIGTL, SLMLP, WSTRP.

Notes: Mycobank 832345. High Ridge Park and Schooley Mill Park specimens identified by examination of spores by R. Solem. Cap turns cherry-/orange-red w/ KOH or NH₄OH. Flesh negative w/ both.

Synonym: Boletus longicurvipes Snell & A.H. Smith

References: BBF 341, 365. BBH 232. BRB1 125. BRB2 112 BBH 232, BRBD 217, 51. K&M 116. Phi 260. Roo 328. W 235-6.



SPORES (13.9-16.0 x 4.7-5.5 μm): Schooley Mill Park 8/1/2018, R. Solem.