

Hairy Panus
Ruddy:Panus

Lentinus strigosus



OVERALL: MPENA, 5/17/2010, J. Solem.

GILLS: MPENA, 5/17/2010, J. Solem.

SPORES (5.2-6.6 X 2.8-3.3 μ m): MPENA, 5/17/2010, R. Solem.

Hairy Panus
Ruddy Panus

[*Lentinus strigosus*](#)
Polyporaceae

J	F	M	A	M	J	J	A	S	O	N	D
			+	+	+	+	+	+	+	+	+

ID: Densely hairy, pinkish cap; fan-shaped. Bitter taste.

Habitat: Solitary, clusters. On hardwoods.

Cap: 1 – 3" [2.5 – 7.5 cm]

Pinkish-tan/reddish-brown (purple tints when young, paler in age).
Densely hairy, velvety/fuzzy. Fan-/funnel-shaped, sometimes lobed.
Margin inrolled and hairy. Flesh white, thin, tough. Taste bitter.

Gills: White to tan. Close, narrow.

Spores: White/off-white toward cream. Elliptical, smooth.

Stalk: Pinkish-tan/light reddish-brown. Short, stubby, mostly lateral, hairy.

Frequency: Occasional.

Locations: MPENA, MTPLT, OTHER, WINTR.

Synonyms: *Lentinellus strigosus*. *Panus rudis*.

Notes: Mycobank 149820. MPENA specimen verified by examination of spores by R. Solem.

References: BBF 184, 297. Bin 367. Lin 791, 505. M&M 141.
Roo 144.