

None

Mallocybe unicolor (Peck) Matheny & Esteve-Rav



TOP: Cedar Lane Park, 6/25/2020,
J. Solem.
 SIDE: Cedar Lane Park, 6/25/2020,
J. Solem.
 GILLS and STALK: High Ridge Park,
 8/15/2020, *J. Solem.*
 SIDE: Donleigh, 8/17/2020,
J. Solem.

None

Mallocybe unicolor
Inocybaceae

J	F	M	A	M	J	J	A	S	O	N	D
				---	---	+	---	+	---	---	

ID: Shaggy dull brown cap and stalk. Use KOH.

Habitat: Scattered, groups. Under hardwoods, especially grassy, urban areas.

Cap: 0.8 – 2" [2 – 5 cm]

Dull orangish-brown. Fibrillose, becoming finely scaly. Convex (nearly flat in age). Flesh whitish. Odor somewhat foul. KOH turn cap dark brown.

Gills: Dull yellow (yellowish-brown in age w/ whitish edges). Attached, close, secondary gills.

Spores: *Medium brown.* Ellipsoid, smooth, hyaline.

Stalk: 0.8 – 2" x 1/8 – 1/4" [2 – 5 cm x 3 – 5 mm]

Brownish, fibrillose-scaly base or overall. Basal mycelium white.


Frequency: Occasional.

Locations: CEDLP, HIRIP, OTHER.

Notes: Mycobank 830477. Cedar Lane Park, Donleigh, and High Ridge Park specimens verified by examination of spores by R. Solem.

Synonym: *Inocybe unicolor.*

References: Kuo. *The Mycophile* May-June 2014, pp 12-13.



SPORES (9.2-10.9 x 4.8-5.2 μm): Cedar Lane Park, 6/25/2020, *R. Solem.*