

Orange-yellow Marasmius

Marasmius strictipes
(Peck) Singer



OVERALL: Woodstock, PVSP, 10/15/2016,
R. Orr.
CAP: Morning Choice Trail, PVSP,
10/13/2015, *J. Solem.*
GILLS and STALK: Morning Choice Trail,
PVSP, 10/13/2015, *J. Solem.*
SPORES (6.7-8.3 x 3.3-3.6 μm): Murray
Hill, 10/13/2015, *R. Solem.*

Orange-yellow Marasmius

[*Marasmius strictipes*](#)
Marasmiaceae

J	F	M	A	M	J	J	A	S	O	N	D
					-+	- -	- - +	- - +	++	- -	

ID: Light yellow to tan, convex, margin uplifted in age. Gills white/pale orangish-yellow.

Habitat: Scattered, groups. Decaying deciduous leaves.

Cap: 0.8 – 2.5 “ [2 – 6.5 cm]

Light yellow/orangish in spots at edge (tan in age). Moist. Convex/broadly umbonate (especially in age). Margin uplifted in age. Flesh white/yellowish-white. Taste mild to peppery.

Gills: Yellowish-white/pale orangish-yellow. Attached but often receding. Narrow. Close/crowded. Secondary gills.

Spores: *White/cream/yellowish.* Ellipsoid, smooth, hyaline.

Stalk: 1 – 3.5” x 1/8 – 3/8” [2.5 – 8.5 cm x 3 – 9 mm]

White/pale yellow. Hollow. Pale yellow/white mycelium binds to mat of leaves. No rhizomorphs.

Frequency: Occasional.

Locations: MCHPV, MURHL, WINTR, WSKPV.

Notes: Mycobank 300226. Species determination Morning Choice, PVSP, specimen by M. Kuo (*pers.com.*). Murray Hill and Samantha Way spores examined by R. Solem.

Synonyms: *Collybia strictipes.* *Gymnopus strictipes.*

References” BBF 202, 301. BBH 133. Bni 186. K&M 274. Kuo. Lin 776, 32. Phi 73. Rog.