## Clustered Bonnet

*Mycena inclinata* (Fries) Qu**é**let



"VERALL, Henryton, PVSP, 10/2/2009, *J. Solem.*TOPS: Morning Choice Trail, PVSP, 10/2/2014, *J. Solem.*SIDE: Morray Hill, 10/9/2020, *J. Solem.*GILLS and STALK: Morning Choice Trail, 10/2/2014, *J. Solem.*SPORES (8.2-10.2 x 5.3-6.9 µm): Murray Hill, 10/9/2020, *R. Solem.*

## Clustered Bonnet <u>Mycena inclinata</u> Mycenaceae J F M A M J J A S O N D

**ID**: Cap gray-brown; bell-shaped. Stalk top light, darker below; white mycelium at base. Odor. May develop yellow stains.

Habitat: Dense clusters. On hardwood (oak) stumps or logs.

## **Cap**: 0.8 – 1.2" [2 –3 cm]

Gray-brown. Conical expanding to bell-shaped, w/ prominent darker umbo (flat in age). Margin slightly overhangs gills, striate. Flesh whitish, thin. Odor rancid/mealy. Taste mild. Fringe-like "teeth" when young.

**Gills**: Whitish/fleshy-pink. Attached. Depth = 2X cap flesh.

Spores: White/cream. Ellipsoid, smooth, hyaline..

**Stalk**: 2.4 - 3.9" x 1/16 - 1/8" [6 - 10 cm x 2 - 4 mm] White at top, yellowish in mid section, darker brown-red towards base (which is covered w/ fine white down). Hollow.

Frequency: Very common.

Locations: HENPV, HIRIP, MCHPV, MURHL, NOLAP, OTHER, PIGTL, ROBNC, ROGOR, RKBRP, SLMLP, SYRIV, WSKPV, WSTRP.

**Notes**: Mycobank 194409. Morning Choice Trail, Murray Hill, and Pigtail specimens identified by examination of spores by R. Solem.

**References**: BBF 209key. BBH 134. Bni 200fn. Lin 780fn. M&M 171. Phi 94. Rog. Stu 139.