

Rooted Collybia

Oudemansiella furfurecea (Peck)
Zhu L. Yang, G.M. Muell., G. Kost & Rexer,



GILLS and STALK: Morning Choice Trail, PVSP, 10/18/2011, *J. Solem*.
 OVERALL: MPEA, 8/30/2012, *J. Solem*.
 GILLS, STALK and ROOT: Morning Choice Trail, PVSP, 10/18/2011, *J. Solem*.
 OVERALL: Lake Kittamaquindi, 5/20/2017, *J. Solem*.
 SPORES (15.6-17.7 X 10.0-11.7 µm): Lake Kittamaquindi, 5/20/2017,
R. Solem.

Rooted Collybia

Oudemansiella furfurecea
Physalacriaceae

J	F	M	A	M	J	J	A	S	O	N	D
				-++	+++	+++	+++	+++	+++	+-	+

ID: Cap brown w/ darker grooved margin; flat (depressed in age), distinct umbo, rubbery skin. Stalk long w/ long underground portion.

Habitat: Solitary, scattered, groups. On ground in mixed forests; infrequently on decayed hardwood logs.

Cap: 0.8 – 4” [2 – 10 cm]
Light brown/dark grayish-brown w/ darker margin. Dry/slippery. Finely velvety (nearly smooth in age). Convex/nearly flat w/ rubbery cuticle; often wrinkled around distinct umbo, grooved (depressed in age). Flesh white.

Gills: White. Attached but typically separating from cap in age. Subdistant. Fairly broad. Secondary gills.

Spores: *White*. Smooth, broadly ellipsoid/fusiform.

Stalk: Above ground portion 3 – 8” x 0.1 – 0.8”
[7.5 – 20 cm x 3 - 2- mm] White beneath grayish/brownish scales/fibers. Usually tapered upward, twisted. Flesh white. Hollow.

Frequency: Abundant.

Locations: BIGBR, BOLPL, CEDLP, DANPV, FRANF, HAMVL, HIRIP, LONGC, MANWD, MCHPV, MPENA, MURHL, OTHER, PIGTL, RKBRP, ROBNC, ROGOR, SAVPK, SYRIV, TRIRS, UMDCF, WINTR, WSTRP.

Notes: Mycobank 512754. Woehrel & Light (338) have key to other species. Lake Kittamaquindi and Rockburn Branch Park records verified by examination of spores by *R. Solem*.

Synonym: *Hymenopellis furfuracea* (Peck)
R.H. Petersen Xerula furfuracea.

References: Bar 303. BBF 270, 314. Bni 172.K&M 394. Kuo. M&M 174 fn. M&S 96. Pac 38. Roo 265. Stu 160. W&L 336-8.