

Belted Panaeolus
Girdled Panaeolus

Panaeolus cinctulus
(Bolton) Saccardo



TOP: Long Corner, PRSP, 7/16/2010, J. Solem.
OVERALL: Long Corner, PRSP, 7/16/2010, J. Solem.
GILLS and STALK: Long Corner, PRSP, 7/16/2010, J. Solem.
OLD SPECIMENS: Long Corner, PRSP, 7/16/2010, J. Solem.

Belted Panaeolus
Girdled Panaeolus

Panaeolus cinctulus
Psathyrellaceae

J	F	M	A	M	J	J	A	S	O	N	D
					---	+	---	---	---		

ID: Brown/black; convex; margin w/ dark zone. Gills brown/black, mottled; edges whitish. Stalk coated w/ white fibrils.

Habitat: Scattered, groups. On horse and cow dung.

Cap: 1.2 – 2" [3 – 5 cm]

Cinnamon-brown. Hygrophanous. Smooth/ wrinkled. Convex/bell-shaped. Broad dark zone around margin. Flesh brownish.

Gills: Mottled dingy brown/black; edges whitish. Attached. Close.

Spores: Black.

Stalk: 1.6 – 4" x 1/8 – 1/4" [4 – 10 cm x 3 – 5 mm]

Reddish-brown/whitish Coated w/ tiny white fibrils. Equal (or both ends slight taper). Hollow.

Frequency: Uncommon.

Locations: LONGC.

Notes: Mycobank 170056. Identification by Lance Biechele (*pers.comm.*)

Synonym: *Panaeolus subbalteatus*

References: Arora 358-359. BBF 217key. Lin 604, 91.

http://en.wikipedia.org/wiki/Panaeolus_cinctulus.