

Hairy Panus  
Ruddy:Panus

*Panus neostrigosus*  
Drechsler-Santos & Wartchow



OVERALL: MPENA, 5/17/2010, J. Solem.  
GILLS: MPENA, 5/17/2010, J. Solem.  
SPORES (5.2-6.6 X 2.8-3.3 µm): MPENA, 5/17/2010, R. Solem.

Hairy Panus  
Ruddy Panus

[\*Panus neostrigosus\*](#)  
Polyporaceae

J	F	M	A	M	J	J	A	S	O	N	D
		+	+++	+++	+++	---	+-	---	+-	---	

**ID:** Densely hairy, pinkish cap; fan-shaped. Bitter taste.

**Habitat:** Solitary, clusters. On hardwoods.

**Cap:** 1 – 3" [2.5 – 7.5 cm]

Pinkish-tan/reddish-brown (purple tints when young, paler in age).  
Densely hairy, velvety/fuzzy. Fan-/funnel-shaped, sometimes lobed.  
Margin inrolled and hairy. Flesh white, thin, tough. Taste bitter.

**Gills:** White to tan. Close, narrow.

**Spores:** White/off-white toward cream. Ellipsoid, smooth.

**Stalk:** Pinkish-tan/light reddish-brown. Short, stubby, mostly lateral, hairy.

**Frequency:** Fairly common.

**Locations:** GUILP, HAMVL, MPENA, MTPLT, OTHER, SLMLP, WINTR.

**Notes:** Mycobank 801445. MPENA specimen identified by examination of spores by R. Solem.

**Synonyms:** *Lentinus strigosus*. *P. rudis*.

**References:** BBF 184, 297. BBH 106. Bin 367. Bni 214.  
BRBD 165, 34. Lin 791, 505. K&M 296. M&M 141 Roo 144. Stu 169.