
Peziza arenaria Osbeck Peziza howsei Roze & Boudier



Peziza arenaria
Peziza howsei
Pezizaceae







OVERALL: Murray Hill, 6/22/2015, *J. Solem.* OVERALL: Murray Hill, 6/22/2015, *J. Solem.* OVERALL: Murray Hill, 6/22/2015, *J. Solem.*

J F M A M J J A S O N D

ID: Body cup-shaped, somewhat violet color.

Habitat: Solitary, clustered on bare soil in forests.

Fruiting Body: 0.8 - 1.2" [2 - 5 cm]

Cup-/disc-shaped Margin smooth/finely scalloped; somewhat inrolled, thin, brittle. Fertile surface: violet/wine-colored (may be barely tinted). Infertile surface: Whitish/ocraceous, slightly tomentose and granular.

Stalk: None. Slightly eccentric attachment to soil, pucker visible on fertile surface


Spores: *Hyaline*. Ellipsoid/fusiform w/ 2 oil drops, patterned.

Frequency: Uncommon.

Locations: OTHER.

Notes: Mycobank 148680 and 225817. R. Solem examined spores of the Murray Hill record. Spores of the above two species differ only in degree of roughness of exterior; that could not be determined. A. E. Bessette concurred in narrowing the choices to the two species listed (*pers. comm.*)

References: BBB 205.



SPORES (16.5-17.3 X 8.7-9.3 μm): Murray Hill, 6/22/2015, R. Solem.