

None

*Pholiota veris* (Singer & A.H. Smith)  
A. H. Smith & Hessler



OVERALL: Gorman Stream Valley NRA, 5/10/2016, *J. Solem*.  
GILLS and STALKS: Gorman Stream Valley NRA, 5/10/2016, *J. Solem*.

None

[\*Pholiota veris\*](#)  
Strophariaceae

J	F	M	A	M	J	J	A	S	O	N	D
			---	+++	---						

**ID:** Yellowish-cinnamon cap, membranous ring (unless old).

**Habitat:** Groups, clusters. On well-decayed deciduous wood/sawdust.

**Cap:** 0.8 – 2.3" [2 – 6 cm].

Pale to yellowish-cinnamon (fades from center outward to pale yellowish-tan in age). Smooth (but moist/sticky/slippery). Margin striate in age. Flesh like cap. Hygrophorous.

**Gills:** Pinkish-tan (cinnamon in age). Covered w/ partial veil at first.

**Spores:** *Brown*. Ellipsoid, w/ or w/o apical pore, smooth, brownish.

**Stalk:** 1.5 – 3.5" x 1/8 – 1.2" [4 – 9 cm x 4 – 12 mm]

Paler than cap. Dry. Scurfy in middle part. Upward-flaring, membranous ring near top (or only zone in age).

**Frequency:** Uncommon.

**Locations:** BOLPL, CEDLP, DANPV, GONRA.

**Notes:** Mycobank 241744. Gorman Stream Valley NRA specimen identified by examination of spores by R. Solem.

**Synonym:** *Kuehneromyces rostratus*

**References:** BBF 230, 306. K&M 311. Kuo. Lin 716fn. M&M 264. Roo 41.



SPORES (6.7-7.4 x 3.7-4.5 μm): Gorman Stream Valley NRA, 5/10/2016, *R. Solem*.