


None

Pluteus aurantiorugosus (Trog) Saccardo



OVERALL: Henryton, PVSP
8/18/2014, *R. Orr*.
GILLS and STALK): Henryton, PVSP.
8/18/2014, *J. Solem*
GILLS and STALK): Henryton, PVSP.
8/18/2014, *J. Solem*
SPORES (6.4 – 6.7 X 4.7 – 4.9 μ m):
Henryton, PVSP. 8/18/2014, *R. Solem*.

None

[*Pluteus aurantiorugosus*](#)
Pluteaceae

J	F	M	A	M	J	J	A	S	O	N	D
						---	++	---			

ID: Brilliant scarlet; broadly convex. White mycelium.

Habitat: Solitary, scattered, groups. Decaying hardwood.

Cap: 3/8 – 3.1" [1 – 8 cm]

Brilliant scarlet/orange-red (orange/yellow in age, darkest over disc). Smooth/slightly velvety, hygrophanous when moist; fibrillose/finely wrinkled in age or dry. Broadly convex (often prominent umbo in age). Margin incurved (flatter when young; striate/finely cracked in age). Flesh yellowish. fragile.

Gills: Whitish (pink in age). Free. Crowded/close. Secondary gills.

Spores: *Pinkish*. Ellipsoid, smooth.

Stalk: 1.1 – 3.5" x 1/8 – 3/8" [3 – 9 cm x 3 – 10 mm]

White/pale yellow (may be orange-red toward base). Striate/fibrillose-scaly (nearly smooth in age). Nearly equal.

Frequency: Uncommon.

Locations: HENPV.

Notes: Mycobank 218699. Henryton, PVSP record identified by examination of spores by R. Solem.

References: Bar 187. BBF 234, 306. Bni 227. K&M 315. Kuo. Lin 675, 230.