

Warted Oak Polypore

Pseudoinonotus dryadeus
(Persoon) T. Wagner & M. Fischer



OVERALL: Daniels, PVSP, 6/10/2015, R. Orr.
OVERALL: Daniels, PVSP, 8/8/2019, R. Orr.
OVERALL: Daniels, PVSP, 6/10/2015, B. Ott.

Warted Oak Polypore

Pseudoinonotus dryadeus
Hymenochaetaceae

J	F	M	A	M	J	J	A	S	O	N	D
			---	+	---	---	++	---	---		

ID: Body whitish-gray (dark in age); uneven/ warty/lumpy. Pores whitish/gray-brown (dark in age). Stalkless. Oozes amber/brown fluid.

Habitat: Solitary, sometimes fused. At base of living oaks or fresh oak stumps; infrequent on other hardwoods.

Fruiting Body: 3 – 16" [7.5 – 40 cm]
Whitish-gray (gray-yellow/red-brown/blackish in age). Hairy (smooth in age), surface uneven/warty, bumpy. Exudes clamber/brown drops when young. Convex/irregular. Flesh (1.5 – 5 cm) yellow-/rusty-brown, fibrous-tough. Odor unpleasant.

Pores: Whitish-/gray-brown (often w/ silvery sheen); dark in age. Round to angular (3 – 5/mm). Tubes (5 – 30 mm).

Stalk: Absent.

Spores: Whitish/pale yellow.

Frequency: Uncommon.

Locations: DANPV, ROGOR.

Notes: Mycobank 474675. See also *Niveoporofomes spraguei*.

Synonyms: *Polyporus dryadeus*. *Inonotus dryadeus*

References: BBH 281. Bin 285. BRBD 259, 61. K&M 224. Lin 466, 534. Rog. W&L 540-1.