

None

Pulveroboletus curtisii (Berkeley) Singer



OVERALL: Hammond Schools, 8/8/2017, *J. Solem*.

OVERALL: Hammond Schools, 8/25/2010, *J. Solem*.

PORES and STALK: Hammond Schools, 8/25/2010, *J. Solem*.

SPORES (10.8-13.0 X 5.1-6.5 μ m): Hammond Schools, 8/25/2010, *R. Solem*.

None

Pulveroboletus curtisii
Boletaceae

J	F	M	A	M	J	J	A	S	O	N	D
							+++	+-	-	-	-

ID: Yellow-orange; convex; viscid.

Habitat: Solitary, scattered, groups. On ground in coniferous or mixed woods.

Cap: 1.1 – 3.5" [3 – 9.5 cm]

Yellow-orange (brownish/whitish areas in age). Convex (broadly convex/flat in age). Viscid/glutinous. Margin w/ narrow band of tissue; incurved when young. Cap stains fingers yellow.

Pores: Whitish/buff/pale yellow (darker in age). Circular/angular (2 – 3/mm). Tubes (6 – 12 mm).

Spores: Olive-brown. Ellipsoid, smooth.

Stalk: 2.4 – 4.8" x 1/4 – 1/2" [6 – 12 cm x 6 – 13 mm]

Pale yellow/yellow. Viscid/glutinous. Scurfy at apex, smooth below. Base w/ white mycelium.

Frequency: Occasional.

Locations: HAMVL, TRIRS.

Notes: Mycobank 290318. *Index Fungorum* only has *Boletus curtisii*. Hammond Schools specimen identified by examination of spores by R. Solem.

Synonym: *Boletus curtisii*

References: BBH 242. BRB1 106, 289. BRB2 294. BRBD 212, 49. K&M 115. M&M 389. W&L 318.