

None

Resupinatus violaceofulvus (Fries) Murrill



TOPS: Bollman Place, 6/20/2014, *J. Solem*.

GILLS: Bollman Place, 6/20/2014, *J. Solem*.

SPORES (6.0 – 7.0 X 2.6 – 3.7 μm): Bollman Place, 6/20/2014, *R. Solem*.

None

[*Resupinatus violaceofulvus*](#)

Tricholomataceae

J	F	M	A	M	J	J	A	S	O	N	D
					++	--	+-	--	--		

ID: Cap dark violet. Gills violet.

Habitat: Solitary, scattered, groups. On dead/decaying conifers.

Cap: 0.2 – 0.4" [3 – 15 mm]

Deep violet-brown. Dry. Coated w/ tiny fibers. Kidney-shaped (from lateral point). Flesh thin, violet-brown.

Gills: Pale violet/violet-brown. Attached. Subdistant. Secondary gills.

Spores: *White*. Sausage-shaped or cylindrical, smooth.

Stalk: Short, stubby, violet-brown w/. fine white hairs..

Frequency: Uncommon.

Locations: OTHER, PIGTL, WINTR.

Notes: Mycobank 507000. Bollman Place and Pigtail records verified by examination of spores by R. Solem.

References: BBF 219, 304.